

AMIGARD

Dosage form: Spray dip for cattle.

Composition: Amitraz 12,5%.

Indications: Controls ticks. Kills lice and mange mites on cattle.

Warnings:

Do not slaughter animals for human consumption with-in 7 days of the last dipping.

Do not use on horses.

Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a wide variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions: Once used the container must be tightly resealed. Destroy empty containers.

Directions for use: Use only as directed. For external use on cattle only. Only use for the spraying of cattle (hand spraying or spray races). Do not mix **Amigard** with any other dip. Fresh dip wash should be prepared for every spraying.

Dosage and mixing instructions: The required amount of **Amigard** should be mixed with water in a suitable container. When it is thoroughly mixed it should be added to the measured bulk of water in the spray sump. This mixture should then be thoroughly stirred. Use one part **Amigard** to 500 parts water as shown in the table below.

Amigard	Water
5 L	2 500 L
1 L	500 L
500 ml	50 L

No more than 500 head of cattle should be sprayed in the spray race at one time, particularly if fouling is a problem. If more than 500 head are sprayed the dip should be boosted at a rate of 100 ml per 100 head of cattle. This ensures that sufficient levels of the active ingredient are maintained on the cattle especially where tick borne diseases are a problem.

Frequency of use:

Intensive tick control: If used as directed at weekly intervals intensive control of ticks is achieved. Ensure that all the animals are treated. Any interruption of the programme should be avoided since the animals will become susceptible to tick borne diseases, particularly those animals that have not been vaccinated.

Mange: Spray at weekly intervals.

Extended treatment intervals: If there is a low tick challenge, the user can at his own discretion spray the animals at intervals of greater than 7 days. It is essential to ensure that all the cattle are vaccinated against the prevalent tick-borne disease.

Presentation: 500 ml, 1 L, 5 L and 20 L.

Storage instructions: Store at room temperature below 25°C. Protect product from extreme heat and damp.

Registration number: G3512 (Act 36/1947).