

For Animal use only

CattleMaster 4

Reg. No. G 2016 Act 36/1947

Namibia Reg. no. NSR 1061

INDICATIONS

CattleMaster 4 is used for vaccination of healthy cattle, including pregnant cows as an aid in preventing infections by infectious bovine rhinotracheitis (IBR) virus, bovine viral diarrhoea (BVD) virus, parainfluenza₃ (PI₃) virus and bovine respiratory syncytial virus (BRSV).

STORAGE INSTRUCTIONS

Store between 2 ° C to 7 ° C. Do not freeze. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency

COMPOSITION

CattleMaster 4 is a freeze-dried vaccine of chemically altered strains of infectious bovine rhinotracheitis (IBR) virus and parainfluenza₃ (PI₃) virus, and modified live bovine respiratory syncytial virus (BRSV), plus a liquid, adjuvant preparation of inactivated cytopathic and non-cytopathic bovine viral diarrhoea (BVD) virus. The liquid component is used to rehydrate the freeze-dried component. Virus fractions are propagated on an established cell line. The product is adjuvanted with aluminium hydroxide to enhance the immune response.

WARNINGS:

Do not vaccinate within 21 days before slaughter.

Keep out of reach of children, uninformed persons and animals.

As with most vaccines, anaphylaxis may occur after use. Initial antidote of adrenalin is recommended and should be followed with appropriate supportive therapy.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitised, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.

Although this vaccine has been tested extensively under a large variety of conditions failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

Use entire contents when first opened.

Sterilized syringes and needles should be used to administer this vaccine. Do not sterilize with chemicals because traces of disinfectant may inactivate the vaccine.

Burn containers and all unused contents.

Contains gentamycin as preservative.

DOSAGE AND DIRECTIONS FOR USE:

General Directions: Vaccination of healthy cattle is recommended. Aseptically rehydrate the freeze-dried vaccine with the diluent provided, shake well and administer 2 ml intramuscularly.

CattleMaster 4 should be administered in the muscular region of the neck.

Primary Vaccination: Administer a single 2 ml dose to healthy cattle, followed by a second dose of **CattleMaster 4** 3 - 4 weeks later. To avoid possible maternal antibody interference with activated immunization, calves vaccinated before the age of 6 months should be revaccinated after 6 months of age.

Revaccination: Annual revaccination with a single dose is recommended.

PRESENTATION

25 by 2 ml doses in two vials

REGISTRATION HOLDER

Pfizer Laboratories (Pty) Ltd

(Reg. No. 1954/000781/07)

85 Bute Lane, Sandton, 2196

P O Box 783720, Sandton, 2146

South Africa

For more information phone 011 320 6000