For Animal Use Only



FELOCELL® 3

Reg. No. G3131 (Act 36/1947)

Only for use by persons authorised thereto in terms of the Veterinary and Para-Veterinary Professions Act, 1982 (Act 19/1982)

INDICATIONS

Felocell 3 is intended for the vaccination of healthy cats and kittens as an aid in preventing feline viral rhinotracheitis (FVR) caused by feline herpesvirus-1, feline respiratory disease caused by feline calicivirus (FCV), and feline panleukopenia (FPL) caused by feline panleukopenia virus.

STORAGE INSTRUCTIONS

Store between 2 °C and 7 °C. Do not freeze.

COMPOSITION

Felocell 3 contains attenuated live virus strains of feline herpesvirus type 1, calicivirus, and panleukopenia virus (Johnson Snow Leopard strain), propagated on established cell lines, and lyophilised to preserve stability.

WARNINGS

Keep out of reach of children, uninformed persons and animals.
Vaccination of pregnant queens should be avoided.

Postvaccination reactions attributable to the vaccine may include lethargy, anorexia, ocular discharge and fever.
If anaphylaxis or hypersensitivity occurs following use, administer adrenaline or equivalent. Without such treatment, serious reactions may result in death.
This vaccine has been shown to be efficacious but some cats may be unable to develop or maintain an adequate immune response following vaccination if they are, for example, incubating any infectious disease, are malnourished or parasitised, or are stressed due to transport or adverse environmental conditions, or otherwise immunocompromised, or the vaccine is not administered according to label directions.
Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Prolonged exposure to higher temperatures may adversely affect potency.
Use entire contents when first opened.
Sterilised syringes and needles should be used to administer this vaccine. Do not sterilise with chemicals because traces of disinfectant may inactivate the vaccine.
Admixing of the vaccine with any other may render it ineffective.
Burn containers and all unused contents.
Contains gentamicin as preservative.

DOSAGE AND DIRECTIONS FOR USE

Use only as directed. Aseptically rehydrate the freeze-dried vaccine with the sterile diluent provided.
General Directions: Vaccination of healthy cats is recommended. Shake well. Aseptically administer 1 ml subcutaneously or intramuscularly
Primary vaccination: Healthy cats 9 weeks of age or older should receive 2 doses, each administered 3 - 4 weeks apart. Cats vaccinated at less than 9 weeks of age should be revaccinated at 9 weeks of age.
Revaccination: Annual revaccination with a single dose is recommended.

PRESENTATION

25 x 1 dose vials with 25 vials packed in a clear plastic, flip top container with a package insert and lot number stickers included.

REGISTRATION HOLDER

Zoetis South Africa (Pty) Ltd
Co. Reg. No. 2012/001825/07
P.O. Box 783720, Sandton 2146
South Africa
Tel. (011) 245 3300 or 0860 ZOETIS (0860 963847)
Zoetis Inc, 601 West Cornhusker Hwy, Lincoln, Nebraska 68521, USA

zoetis

Namibia: V02/24.2/923 NS7

Zimbabwe: 86/80.23.16/152 (PP Vet)

10/2016
50264300

Slegs vir Dieregebruik

FELOCELL® 3



Reg. Nr. G3131 (Wet 36/1947)

Slegs vir gebruik deur persone gemagtig daartoe in terme van die Wet op Veterinêre en Para-Veterinêre Beroepe, 1982 (Wet 19/1982)

INDIKASIES

Felocell 3 is bedoel vir die inenting van gesonde katte en klein katjies. Dit is 'n hulpmiddel in die voorkoming van kat virus rinotrageïtis wat veroorsaak word deur kat herpesvirus-1, kat respiratoriese siekte veroorsaak deur kat calicivirus, en kat panleukopenie veroorsaak deur kat panleukopenie virus.

BERGINGSAANWYSINGS

Berg tussen 2 °C en 7 °C. Moet nie bevries nie.

SAMESTELLING

Felocell 3 bevat verswakte lewendige virus stamme van kat herpesvirus tipe 1, calicivirus, en panleukopenie virus (Johnson Snow Leopard stam), vermeerder op 'n stabiele kat sellyn en vriesgedroog om stabiliteit te verseker.

WAARSKUWINGS

Hou buite bereik van kinders, oningeligte persone en diere.
Inenting van dragtige katte moet vermy word.

Reaksies na inenting wat toegeskryf kan word aan die entstof is moontlik en kan lusteloosheid, anoreksie, koors en ooguitloopsel insluit. Indien anafilakse of hipersensitiwiteit na gebruik voorkom, dien adrenalien of ekwivalent toe. Sonder sodanige behandeling mag erge reaksies lei tot die dood.
Hierdie entstof is doeltreffend bewys maar sommige diere mag nie in staat wees om 'n voldoende immuunreaksie te ontwikkel of te onderhou na inenting nie. Dit mag gebeur as hulle bv. enige ander siekte inkubeer, ondervoed, geparasiteer of gestres is deur vervoer of swak omgewingstoestande of immuunonderdruk is of as die entstof nie volgens voorskrif toegedien is nie.
Alhoewel die entstof breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

Langdurige blootstelling aan hoër temperature mag die doeltreffendheid nadelig beïnvloed.
Gebruik die totale inhoud nadat dit oopgemaak is.
Steriele spuite en naalde moet gebruik word om die entstof mee toe te dien. Moet nie met chemikalieë ontsmet nie aangesien oorblyfsels daarvan die entstof mag inaktiveer.
Vermenging van die entstof met enige ander middel of entstof mag dit nadelig beïnvloed.
Verbrand houers en alle ongebruikte inhoud.
Bevat gentamisien as preserveermiddel.

DOSIS EN GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui. Rehidreer die vriesgedroogde komponent asepties met die steriele vloeistof wat voorsien word en skud goed.
Algemene aanbevelings: Inenting van gesonde katte word aanbeveel. Skud goed. Dien 1 ml asepties onderhuids of binnespiers toe.
Primêre inenting: Gesonde katte van 9 weke of ouer moet 2 inentings kry, 3 - 4 weke uitmekaar. Indien katjies jonger as 9 weke ouderdom vir die eerste keer ingeënt word, moet die katjies weer op die ouderdom van 9 weke ingeënt word.
Herinenting: Jaarlikse herinenting met 'n enkele dosis word aanbeveel.

AANBIEDING

25 x 1 dosis flessies met 25 flessies steriele oplosmiddel verpak in 'n helder plastiek, opklap houer met 'n voubiljet en lot nommer plakkers ingesluit.

REGISTRASIEHOUER

Zoetis South Africa (Pty) Ltd
Mpy. Reg. Nr. 2012/001825/07
Posbus 783720, Sandton 2146
Suid-Afrika
Tel. (011) 245 3300 of 0860 ZOETIS (0860 963847)

zoetis

Namibië: V02/24.2/923 NS2

Zimbabwe: 86/80.23.16/152 (PP Vet)

10/2016
50264300