

PARACIDE

Dip for cattle, sheep and goats.

For external animal use only.

Composition:

Alphamethrin 7% m/v.

Indications:

Cattle: Controls ticks. Protects against nuisance flies, biting flies and midges and controls face flies. Kills biting and sucking lice and sand tampans infesting cattle.

Sheep and goats: Controls ticks (including Karoo Paralysis ticks on sheep). Kills biting and sucking lice.

Warnings:

1. Handle with care. Poisonous when swallowed or inhaled.
2. Keep out of reach of uninformed persons, children and pets.
3. Store away from food and feed.
4. Toxic to bees, other insects, fish and wildlife.
5. Allow one day between last treatment and slaughter for human consumption in cattle and 7 days for sheep and goats. Milk from treated cows can be used for human consumption immediately after application.
6. Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions:

1. Do not mix **Paracide** with any other products.
2. Avoid contact with concentrate. Wash immediately with soap and water after accidental skin contact. Contact with the dip mixture may cause a burning sensation of the skin and eyes which may last for 24 hours. It is, however, transient and without harmful effect.
3. Avoid prolonged contact with the spray mixture or inhalation of spray mist.
4. Wash hands thoroughly before eating, drinking or smoking.
5. Prevent contamination of food, utensils, drinking water, rivers, dams, grazing, etc.
6. Destroy empty containers completely by perforation and flattening. Do not re-use for any other purpose.
IN CASE OF ACCIDENTAL POISONING OR INGESTION SEEK IMMEDIATE MEDICAL ATTENTION.
No specific antidote, treat symptomatically.

Dosage and directions for use:

Use only as directed.

Dilution: Spray race, hand spraying and plunge dip for cattle. Belly bath for sheep and plunge dip for goats.

Freshfill: 1 L **Paracide** per 1000 L water.

For plunge dip replenishment for cattle: 1,2 L **Paracide** per 1000 L water.

For constant replenishment for sheep and goats: 1,4 L **Paracide** per 1000 L water.

Spray race: Spray **Paracide** at regular intervals according to tick challenge. It is good spray race practice not to carry over mixture from one spraying to the next and preferably to make up only sufficient mixture for one treatment.

Hand spraying: Prepare about 10 litres of mixture per animal. Wet animals thoroughly, particularly in the ears, under the tail and in the brush of the tail.

Plunge tanks and belly baths: Add the required amount of **Paracide** to an equal quantity of water in a bucket or drum. Mix well by stirring. Add the creamy emulsion to the diptank and mix thoroughly with a suitable stirrer. First chase through 30 animals to ensure proper mixing and return these for redipping.

Lice: Two applications with an interval of one week is recommended for cattle. One application should be sufficient for sheep and goats.

Flies: Spray or dip with **Paracide** at regular intervals according to fly challenge.

Presentation: Containers of 1 litre, 5 litre and 20 litre.

Registration number: G791 (Act 36/1947).