

## FOR ANIMAL USE ONLY/SLEGS VIR DIEREGERBRUIK

### WARNINGS

Keep out of reach of children and uninformed persons and animals. Wash hands after use.

Do not spray in or near the eyes.

No withdrawal period for meat or milk intended for human consumption. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### MODE OF ACTION

Oxytetracycline is a product of the metabolism of *Streptomyces rimosus* and is one of the family of tetracycline antibiotics. Oxytetracycline is primarily bacteriostatic and is thought to exert its antimicrobial effect by inhibition of protein synthesis. It is active against a wide range of gram-negative and gram-positive organisms.

### DOSAGE AND ADMINISTRATION

For topical use only

Shake well before use

Hold can 15-20 cm from the area and spray onto affected area for 1-2 seconds. Apply one or more times a day depending on patient interference or environmental conditions.

### PRECAUTIONS

The container is pressurised, do not expose to heat or sunlight.

Do not puncture or incinerate even when empty.

BATCH NO:

EXPIRY DATE:

LOT NO:

VERVAL DATUM:

## TERRAMYCIN WOUND SPRAY

**Reg. No G 2234 Act/Wet 36/1947**  
**Namibia reg. no NSR 0709**

### INDICATIONS

For the prevention and treatment of topical infections sensitive to oxytetracycline e.g. lacerations, abrasions, wound dehiscence, dermatitis, footrot in cattle, sheep, goats, pigs, horses, dogs and cats

### INDIKASIES

Vir die voorkoming en behandeling van topikale infeksies gevoelig vir oksitetrasiklien bv. skuurwonde, skaafplekke dermatitis en vrotpootjie. van beeste, skape, bokke, varke, perde, honde en katte.

### CAUTION/VERSIGTIG

Store in a dry place below 25 °C

Bêre in 'n koel plek onder 25 °C

**OXYTETRACYCLINE HCl 4g**  
**WITH A BLUE MARKER DYE**  
**OKSITETRASIKLIE HCl 4g**  
**MET BLOU MERKER KLEURSTOF**

**Registration Holder/Registrasie Houer**

PFIZER LABORATORIES

REG. NO 1954/000781/07

Bute Laan 85 Bute Lane

SANDTON

2196

**Net content 150 ml Netto inhoud**

### WAARSKUWINGS

Hou buite die bereik van kinders en oningeligte persone en diere.

Was hande na gebruik.

Moet nie in of om die oë spuit nie.

Geen onttrekkingsperiode is nodig vir vleis of melk wat vir menslike gebruik bestem is nie. Alhoewel hierdie middel onder 'n wye verskeidenheid toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

### WERKING

Oksitetrasiklien is 'n produk van die metabolisme van *Streptomyces rimosus*, en behoort to die tetrasiklien familie van antibiotika. Oksitetrasiklien is primêr bakteriostaties. Dit het 'n antimikrobiële uitwerking deurdat dit proteïensintese inhibeer. Dit is aktief teen 'n groot verskeidenheid gram-negatiewe en gram-positiewe organismes.

### DOSIS EN GEBRUIKSAANWYSINGS

Vir topikale gebruik alleenlik

Skud goed voor gebruik. Hou die spuitkan 15-20 cm vanaf area en spuit op geaffekteerde area vir 1-2 sekondes. Toediening kan 1 of meer keer per dag geskied afhangend van inmenging van die pasient en omgewingstoestand.

### VOORSORGMAATREËLS

Bêre in 'n droë plek benede 25°C.

Die houer is onder druk. Moet nie aan hitte of sonlig blootgestel word nie.

Moet nie prik of brand nie al is dit leeg.

