

FOR ANIMAL USE ONLY

VALBANTEL®

Reg. No. G732 (Act 36/1947)

Indications

Roundworm, Lungworm, Milk Tapeworm, Liver Fluke, Nasal Worm remedy for Sheep and Goats with residual protection against re-infestation of Wireworm and Hookworm.

EFFICACY:

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

1. Roundworm efficacy:

Species	Immatures	Adults
Wireworm	*	*
Brown Stomachworm	*	*
Bankruptworm	*	*
Long-necked Bankruptworm	*	*
Hookworm	*	*
Nodular worm	*	*
Large-mouthed bowelworm	*	*
Lungworm	*	*

Definition: * Control (≥ 90 % effective)

2. Residual efficacy:

Interval between treatment and reinfestation	Wireworm	Hookworm
3 weeks	*	*

Definition: * Control (≥ 90 % effective)

3. Tapeworm efficacy:

Milk tapeworm (<i>Moniezia</i> spp.)	Class 1
---------------------------------------	---------

Definition: Class 1: 100 % effective in more than 80 % of the treated flock.

4. Fluke efficacy:

Fluke species	Stage	
	6 weeks	Adults
Liver Fluke (<i>Fasciola hepatica</i>)	**	*
Giant Liver Fluke (<i>F.gigantica</i>)	**	*

Definition: * Control (≥ 90 % effective)

** Aid in Control (60 - 89 % effective)

5. Nasal worm efficacy: (*Oestrus ovis*)

Control (≥ 90 % effective) 1st, 2nd and 3rd instar larvae.

6. **Ovicidal:** Prevents roundworm eggs present in the animal at dosing from hatching.

PRODUCT

Contains: Albendazole 1,9 % m/v and Closantyl sodium 3 % m/v. *Benzimidazoles & Salicylanilides/Substituted Phenols

PRECAUTIONS

Poisonous; Store in a cool place.

WARNINGS:

ALLOW 30 DAYS BETWEEN LAST DOSING AND SLAUGHTER FOR HUMAN CONSUMPTION.

NOT FOR USE IN MILK GOATS WHOSE MILK IS INTENDED FOR HUMAN CONSUMPTION.

Handle with care.

Keep out of reach of children, uninformed persons and animals.

Management practices should always ensure that all drenching procedures are carefully carried out.

It is particularly important not to exceed the recommended dosage in ewes during mating or during the first month of pregnancy.

Prevent contamination of food, food utensils, etc.

Wash hands thoroughly after use.

Do not repeat treatments more frequently than every 3 weeks.

It is recommended that all animals be vaccinated against pulpy kidney disease before any worm remedy is dosed.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DOSAGE AND DIRECTIONS FOR USE: Use only as directed. Shake well before use.

DOSAGE: 2 ml per 10 kg livemass, eg:

Mass	20 kg	30 kg	40 kg	50 kg
Dosage	4 ml	6 ml	8 ml	10 ml
No. of doses per pack	125	83	62	50

REGISTRATION HOLDER:

Pfizer Laboratories (Pty) Ltd

85 Bute Lane

SANDTON

2196

Tel. (011) 320 6000