

FOR ANIMAL USE ONLY

VALBAZEN® ULTRA

Reg. No. G2037 (Act 36/1947)

Roundworm, Lungworm and Milk Tapeworm remedy for Cattle, Sheep and Goats, and Liver Fluke remedy for Sheep and Goats.

EFFICACY:

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

1. Roundworm efficacy:

Species	Immatures	Adults
Wireworm	*	*
Brown Stomachworm	*	*
Cattle Bankruptworm	*	*
Hookworm	*	*
Nodular Worm	*	*
Lungworm	*	*

Definition: * Control (≥ 90 % effective)

2. Tapeworm efficacy:

Milk tapeworm (<i>Moniezia</i> spp.)	Class 1
---------------------------------------	---------

Definition: Class 1: 100 % effective in more than 80 % of the treated herd.

3. Ovicidal: Prevents roundworm eggs present in the animal at dosing from hatching.

SHEEP AND GOATS: Dosage: Dilute 1 part (eg. 2 litre) Valbazen Ultra with 3 parts (eg. 6 litre) of clean water and dose at a rate of 2 ml per 10 kg live mass, eg.

Mass	20 kg	30 kg	40 kg	50 kg
Dosage	4 ml	6 ml	8 ml	10 ml
No. of doses per pack	2000	1333	1000	800

EFFICACY:

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

1. Roundworm efficacy:

Species	Immatures	Adults
Wireworm	*	*
Brown Stomachworm	*	*
Bankruptworm	*	*
Long-neck bankruptworm	*	*
Hookworm	*	*
Nodular worm	*	*

Large-mouthed bowelworm	*	*
Lungworm	*	*

Definition: * Control (≥ 90 % effective)

2. Milk tapeworm efficacy:

Milk tapeworm (<i>Moniezia</i> spp.)	Class 1
---------------------------------------	---------

Definition: Class 1: 100 % effective in more than 80 % of the treated flock.

3. Liver Fluke efficacy: Aid in control of adult Liver Fluke (*Fasciola hepatica* and *F.gigantica*).

Definition: Aid in control (60 % - 89 % effective)

4. Ovicidal: Prevents roundworm eggs present in the animal at dosing from hatching.

PRODUCT

Contains: Albendazole 7,6 % m/v.

PRECAUTIONS

CAUTION

Store in a cool place.

WARNINGS:

THE MILK OF TREATED ANIMALS MAY NOT BE USED FOR HUMAN CONSUMPTION OR MANUFACTURING PURPOSES OR MIXED WITH NORMAL MILK WITHIN 72 HOURS AFTER DOSING. DO NOT SLAUGHTER CATTLE FOR HUMAN CONSUMPTION WITHIN 12 DAYS AND SHEEP AND GOATS WITHIN 10 DAYS OF DOSING.

Keep out of reach of children, uninformed persons and animals.

The diluted product may not be used after 6 weeks.

Management practices should always ensure that all drenching procedures are carefully carried out. It is particularly important not to exceed the recommended dosage in ewes during mating or during the first month of pregnancy.

It is recommended that all sheep and goats be vaccinated against pulpy kidney disease before any worm remedy is dosed.

Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DOSAGE AND DIRECTIONS FOR USE:

Use only as directed. Shake well before use. For sheep and goats, first dilute 1 part Valbazen Ultra to 3 parts of clean water before use.

CATTLE: Dosage: The dose is 1 ml per 10 kg live mass, eg.

Mass	100 kg	150 kg	200 kg	250 kg	300 kg
Dosage	10 ml	15 ml	20 ml	25 ml	30 ml
No. of doses per pack	200	133	100	80	66

DOSAGE FOR LIVER FLUKE (AFTER DILUTION): 2,5 ml per 10 kg live mass, eg.

Mass	20 kg	30 kg	40 kg	50 kg
Dose	5 ml	7,5 ml	10 ml	12,5 ml
No. of doses per pack	1600	1066	800	640

REGISTRATION HOLDER:
Pfizer Laboratories (Pty) Ltd
Reg. no. 1954/000781/07
85 Bute Lane
SANDTON
2196