

FOR ANIMAL USE ONLY

VibraVet® 10% Paste

Reg. No. G3552 Act 36/1947

Namibia reg. no. NSR 1340

For Cats and Dogs

INDICATIONS

VibraVet* 10% Paste is effective against important pathogens which affect dogs and cats including *Staphylococcus* spp., *Streptococcus* spp., *Escherichia coli*, *Pasteurella* spp., *Klebsiella* spp., *Proteus* spp., *Pseudomonas* spp., *Enterobacter* spp., *Actinobacter* spp., and *Alicalgines* spp., as well as *Mycoplasma* spp. and *Chlamydia* spp. VibraVet* 10% Paste is indicated for the treatment of infections in cats and dogs caused by or associated with doxycycline sensitive organisms including:

- Skin, eg. pyoderma, folliculitis
- Respiratory infections, eg. rhinitis, pharyngitis, bronchitis, pneumonia
- Urogenital infections
- Gastrointestinal tract infections
- Puerperal infections
- Osteomyelitis
- Otitis media

and secondary bacterial infections associated with canine distemper and feline panleukopenia.

STORAGE INSTRUCTIONS

Store below 25°C.

WARNINGS

Do not use in pregnant bitches in the last 3 weeks of pregnancy and in newborn animals during the first month of life. Vomiting may occur as an uncommon side-effect. Keep out of reach of children and unformed persons.

Although this remedy has been extensively tested under a large variety of conditions, failure may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

The use of tetracyclines in young animals prior to the eruption of adult teeth may lead to permanent yellow staining of teeth. Antacids, containing aluminium, calcium and magnesium salts may affect the absorption of doxycycline and therefore should not be given to animals treated with VibraVet. Dispose of empty container by wrapping with paper and placing in rubbish bin.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

For both dogs and cats a loading dose of 5mg doxycycline/kg should be administered orally followed at 12 hourly intervals by two doses of 2.5 mg/kg. Maintenance doses of 2.5 mg/kg are then given as single dose at 24 hour intervals for a further 5 days. The recommendation should be made when dispensing doxycycline that it be administered either in conjunction with or immediately following feeding.

DOSAGE GUIDE:

Adult Dogs/Cats

Body Mass	5 kg	10 kg	20 kg	30 kg	40 kg
Loading dose (g)	0.25	0.5	1.0	1.5	2.0
Follow-up dose (g)	0.125	0.25	0.5	0.75	1.0

Therapy should be continued for 5-7 days or for 48 hours after all symptoms have subsided. In chronic or refractory cases a longer course of therapy may be required. Based on multi-centre trials, the following durations are suggested as guidelines:

Chronic skin disease 10-20 days
Chronic cystitis 10-28 days
Respiratory disease 8-10 days

PACKAGING

Packed in cartons of 20 x 2.5 g oral dosing syringes.

REGISTRATION HOLDER:

Pfizer Laboratories (Pty) Ltd
(Reg. No. 1954/000781/07)
85 Bute Lane, Sandton 2196
PO Box 783720, Sandton 2146
South Africa
Tel: 011-320-6000