

For Animal Use Only

## TrichGuard®

Reg. No. G1749 (Act 36/1947)  
N-SR 1099 (Namibia)

### INDICATIONS

For vaccination of healthy cattle as an aid in the prevention of disease caused by *Tritrichomonas foetus*. Neomycin and polymyxin B added as preservatives.

### DESCRIPTION

TrichGuard® contains killed, concentrated cultures of *Tritrichomonas foetus* suspended in a special oil adjuvant.

### STORAGE INSTRUCTIONS

Keep refrigerated between 2 °C and 7 °C. AVOID FREEZING. Keep in the dark.

### WARNINGS

Do not slaughter cattle for human consumption within **60 days** of treatment.

Keep out of reach of children, uninformed persons and pets.

The vaccine is for prophylactic use only i.e. it will not alter the clinical course of the disease in cattle with pre-existing infection.

Only healthy, immune-competent cattle should be vaccinated.

Should an allergic reaction occur for any reason, intramuscular adrenaline should be administered.

In any cattle population, a small number of individuals may fail to respond fully to vaccination. Development of adequate immunity may also be altered by other conditions such as hereditary factors, subclinical infection, nutrition, disease and parasitism.

Disinfectants will render the vaccine ineffective.

Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons.

If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS

Use entire contents when first opened.

SHAKE WELL before use.

Transient swelling may occur at the injection site.

### DIRECTION FOR USE: Use only as directed.

#### General Directions:

Administer one 2 ml dose subcutaneously in the neck or behind the shoulder using aseptic technique.

A second dose should be administered 2 to 4 weeks later. The last injection should precede the breeding season by not less than 4 weeks.

**Annual Revaccination:** An annual booster vaccination with a single dose should precede the breeding season by not less than 4 weeks.

### PACKAGING

TrichGuard® vaccine for cattle is packed in 50 dose (100 ml) glass or plastic vials.

### SHELF LIFE

The expiry date stated on the packaging is only guaranteed if the vaccine is stored in the dark at a temperature between 2 °C and 7 °C. The expiry date is 24 months from date of manufacture.

### REGISTRATION HOLDER

Pfizer Laboratories (Pty) Ltd  
Reg. No. 1954/000781/07  
P.O. Box 783720, Sandton 2146  
85 Bute Lane, Sandton 2196, South Africa.  
Tel.: (011) 320 6000



® TrichGuard and Pfizer logo are registered trade marks.

Rev. 02/2012

233662/20